

Superatom molecular orbitals: a new type of long-lived electronic states

Y. Pavlyukh* and J. Berakdar

Institut für Physik, Martin-Luther-Universität Halle-Wittenberg, Heinrich-Damerow-Str. 4, 06120 Halle, Germany

(Dated: October 5, 2011)

We present *ab initio* calculations of the quasiparticle decay times in a Buckminsterfullerene based on the many-body perturbation theory. A particularly lucid representation arises when the broadening of the quasiparticle states is plotted in the angular momentum (ℓ) and energy (ϵ) coordinates. In this representation the main spectroscopic features of the fullerene consist of two occupied nearly parabolic bands, and delocalized plane-wave-like unoccupied states with a few long-lived electronic states (the superatom molecular orbitals, SAMOs) embedded in the continuum of Fermi-liquid states. SAMOs have been recently uncovered experimentally by M. Feng, J. Zhao, and H. Petek [Science **320**, 359 (2008)] using scanning tunneling spectroscopy. The present calculations offer an explanation of their unusual stability and unveil their long-lived nature making them good candidates for applications in the molecular electronics. From the fundamental point of view these states illustrate a concept of the Fock-space localization [B. L. Altshuler, Y. Gefen, A. Kamenev, and L. S. Levitov, Phys. Rev. Lett. **78**, 2803 (1997)] with properties drastically different from the Fermi-liquid excitations.

PACS numbers: 71.10.-w, 31.15.A-, 71.20.Tx, 71.10.Ay, 73.22.Dj

Superatom molecular orbitals (SAMOs) were recently discovered¹ as universal characteristics of C_{60} molecules and their aggregates. Further experiments on endohedral systems² and calculations for a series of quasi-spherical molecules³ showed that SAMOs are associated with the whole system (rather than with a particular atom), have a well defined spherical symmetry (Fig. 1, angular momentum $\ell = 0, 1, 2$, principal quantum number $n = 3$) and are capable of forming chemical bonds. Being markedly different from other unoccupied delocalized states they hold a promise for unique applications in molecular electronics. Hitherto theory was not able to answer why these states are so resilient to the chemical environment and why their character is not washed out by the hybridization: questions of critical importance for the SAMO-mediated charge transport⁴.

To resolve these issues we report here results of quantum chemical calculations for prototypical C_{60} using recently developed self-energy formalism⁵. When a transport electron is injected into the unoccupied SAMO (the single-particle ($1p$) state α) the molecule undergoes a transition into the mixed quantum state that decays in time. At later instances the decay is exponential $\exp(-\gamma_\alpha t)$ with γ_α given by the imaginary part of the on-shell self-energy $\gamma_\alpha = \Im \Sigma_{\alpha\alpha}(\epsilon)$ (Ref. 6). However, it is the initial stage of the decay that is manifested in ultrafast nonequilibrium processes of technological relevance. We have shown⁵ that the spectral function for a large class of relevant electronic systems can be represented in the form

$$\mathbf{A}_\alpha(t) = \exp\left(-\gamma_\alpha \frac{t^2}{t + \tau_\alpha}\right), \quad (1)$$

where the set-in time of the exponential decay is given by $\tau_\alpha = 2\gamma_\alpha/\sigma_{\alpha\alpha}^2$. The spectral function $\mathbf{A}_\alpha(t)$ has the following short and long time-limits:

$$\frac{d}{dt}\mathbf{A}_\alpha(t) \xrightarrow{t \rightarrow 0} -\sigma_{\alpha\alpha}^2 t, \quad (2)$$

$$\mathbf{A}_\alpha(t) \xrightarrow{t \rightarrow \infty} e^{-\gamma_\alpha t}, \quad (3)$$

meaning that for short times ($t \ll \tau_\alpha$) we have the quadratic




FIG. 1. Single-particle states of C_{60} molecule. Each group of the degenerate states is represented by a cylinder and is characterized by the averaged orbital momentum (x -axis), the energy (y -axis), the energy uncertainty σ (atomic units (a.u.), z -axis or colour coding) and the orbital momentum uncertainty ($\delta\ell$, the area of the cylinder base). The occupied valence states (as calculated fully *ab-initio*) form two nearly parabolic bands that can be described analytically by $E_n(\ell) = E_n(0) + \frac{\ell(\ell+1)}{2R^2}$, $n = 1, 2$. This is a signature of the delocalized motion of the electrons on the spherical cage of the radius $R = 3.57 \text{ \AA}$ (Ref. 3). SAMOs show comparable values of $\delta\ell$ and much smaller σ and form a third band dissolving in the sea of other unoccupied states. Note that the weak dispersion of the SAMOs with ℓ , as observed above, is due to their larger averaged radius R , as compared to the first two angular subbands.




FIG. 2. Diagrams for the electron self-energy: The exact expression in terms of the $2p1h$ response function ($G^{(3)}$) in which the entrance and the exit channels cannot be separated by cutting one fermion line can be factorized approximately into the exact particle-hole ($p-h$) propagator (Π) and the exact $1p$ Green's function (G) or, equivalently, into a product of the exact screened Coulomb interaction (W) and the exact Green's function.

decay

$$\mathbf{A}_\alpha(t) \sim 1 - \sigma_{\alpha\alpha}^2 t^2 / 2. \quad (4)$$

$\sigma_{\alpha\alpha}^2$ is the central quantity for our theory. It can be determined as follows. A simple substitution of asymptotic expansions into the Dyson equation leads to the exact relations between the spectral moments of the self-energy $\Sigma(\omega)$ and of the spectral function $\mathbf{A}(\omega)$ (Ref. 7):

$$\mathbf{M}^{(0)} = \mathbf{I}, \quad \Sigma_\infty = \mathbf{M}^{(1)} - \varepsilon, \quad (5)$$

$$\Sigma^{(0)} = \mathbf{M}^{(2)} - [\mathbf{M}^{(1)}]^2, \quad (6)$$

where Σ_∞ is the frequency independent real part of the self-energy⁸. ε is a diagonal matrix with the elements given by the

zeroth-order state energies. By defining the matrix of spectral functions in terms of the imaginary part of the single-particle Green function ($\mathbf{A}(\omega) = \frac{1}{\pi} |\text{Im} \mathbf{G}(\omega)|$) and likewise for the spectral function of the self-energy ($\mathbf{S}(\omega) = \frac{1}{\pi} |\text{Im} \Sigma(\omega)|$), and by the use of the superconvergence theorem⁹ one can redefine the matrices in terms of the frequency integrals:

$$\mathbf{M}^{(n)} = \int_{-\infty}^{\infty} d\omega \omega^n \mathbf{A}(\omega), \quad n = 0 \dots 2, \quad (7)$$

$$\Sigma^{(0)} = \int_{-\infty}^{\infty} d\omega \mathbf{S}(\omega). \quad (8)$$

Equation (6) expressed in the basis of Hartree-Fock states is central for our discussion. In order to be feasible it should be written in a finite basis and coupled with an appropriate approximation for the self-energy operator. It is well known that the self-energy can be recast in terms of the $2p1h$ ($2h1p$) six-point correlation function^{8,10}. Our approach corresponds to its factorization into the product of $p-h$ and $p(h)$ correlation functions (Fig. 2) linking it to the established GW approximation^{11,12}. It is remarkable that after all complicated Lehmann-type expressions for the correlation functions are substituted into the $\Sigma = i\mathbf{G}\mathbf{W}$ and the frequency integral in Eq. (8) is performed we obtain an expression for the energy-uncertainty of a particularly simple form:

$$\sigma_{\alpha\beta}^2 = \mathbf{M}_{\alpha\beta}^{(2)} - [\mathbf{M}^{(1)}]_{\alpha\beta}^2 = \Sigma_{\alpha\beta}^{(0)} = 2 \sum_i \sum_{j,k,k'} \langle \alpha i | j k \rangle \langle \beta i | j' k' \rangle [n_j n_{j'} \delta_{kj} \delta_{k'j'} + n_k (1 - n_j) \delta_{kk'} \delta_{jj'} - \rho_{k'j'} \rho_{kj}], \quad (9)$$

where $\rho_{ij} = \langle 0 | c_i^\dagger c_j | 0 \rangle$ is the single-particle density matrix and the Coulomb matrix elements are defined as $\langle \alpha \beta | \gamma \delta \rangle = \int d(\mathbf{r}_1 \mathbf{r}_2) \phi_\alpha(\mathbf{r}_1) \phi_\beta(\mathbf{r}_1) \phi_\gamma(\mathbf{r}_2) \phi_\delta(\mathbf{r}_2) / |\mathbf{r}_1 - \mathbf{r}_2|$. The prefactor 2 arises from the spin degeneracy. Apart from the factorization of the six-point correlation function Eq. (9) is exact. It is sufficient to determine only the density matrix of the correlated ground state. For small molecules or clusters this can be done at any desired level of accuracy by the configuration interaction (CI) approach which, however, currently not feasible for fullerenes. Therefore, we consider a further approximation widely known as GW^0 (Ref. 13) and treat the screened Coulomb interaction on the random phase approximation level. This is equivalent to the configuration interaction singles (CIS) treatment of the excited states. In view of the Brillouin's theorem¹⁴ the ground state density matrix is $\rho_{ij} = n_i \delta_{ij}$ leading to the energy-uncertainty expressible in terms of the Coulomb matrix elements only:

$$\sigma_{\alpha\beta}^2 = 2 \sum_i \sum_{j,k} \langle \alpha i | j k \rangle \langle \beta i | j k \rangle n_k (1 - n_j). \quad (10)$$

The main computational burden is the transformation of the Coulomb matrix elements from the atomic to molecular orbital basis which scales as $\mathcal{O}(N^5)$ in the brute force imple-

mentation. To obtain convergent results we used very large basis sets (up to 6-311++G(3df,3pd)) resulting in $N = 2340$ functions for C_{60} . Integral transformations for this number of basis functions is not feasible with any standard quantum chemistry package and required a parallelized implementation fully accounting for the symmetry of the system.

Our calculations clearly indicate a peculiarity of SAMOs: a strong localization in the energy domain (cf. $\sigma_{\text{HOMO}} = 16.0$ eV and $\sigma_{\text{SAMOd}} = 6.1$ eV) (Fig. 1) or their extended life-time even in comparison with the life-time of HOMO or LUMO states. This finding endorses the potential of SAMOs as transport channels in molecular electronic devices, since the energy is hardly dissipated during the short transport time. Our results are also unexpected from the Landau's theory of Fermi liquids¹⁵ and illustrate that finite systems possess electronic excitations that differ drastically from quasiparticles in extended matter. Namely, a particle excitation may decay into two particles and one hole. This process may recur for many generations or it may stop after a few. For the former case a large number of many-particle peaks form a Lorentzian envelope, the so-called *quasiparticle*. If the decay stops after a finite number of branchings, only Dirac peaks appear in the single particle spectrum (denoting a localization in the Fock

space). A decisive discriminating factor was shown to be the particle's energy (ϵ) (Ref. 16). Our *ab initio* approach suggests that it is actually the kinetic energy ϵ_K that governs the decay. We find that even though the energies of electrons with different radial character may be very similar their kinetic energy is substantially different (as follows from the specific form of the Kohn-Sham potential for this system, Ref. 17), which imposes strong restrictions on the Coulomb matrix elements and

leads to the localization of SAMOs in the Fock space. This hints on the prolonged life-times and stipulates the observed high stability of these states.

We thank the DFG for financial support through SFB 762 and H. Petek for fruitful discussions. J.B. acknowledges financial support by Stanford Institute for Materials & Energy Science and Stanford Pulse Institute for ultrafast science where parts of this work were completed.

* yaroslav.pavlyukh@physik.uni-halle.de

¹ M. Feng, J. Zhao, and H. Petek, *Science*, **320**, 359 (2008).

² J. Zhao, M. Feng, J. Yang, and H. Petek, *ACS Nano*, **3**, 853 (2009).

³ Y. Pavlyukh and J. Berakdar, *Chem. Phys. Lett.*, **468**, 313 (2009).

⁴ T. Huang, J. Zhao, M. Feng, H. Petek, S. Yang, and L. Dunsch, *Phys. Rev. B*, **81**, 085434 (2010).

⁵ Y. Pavlyukh, A. Rubio, and J. Berakdar, arXiv:1107.5632v1.

⁶ Y. Pavlyukh, J. Berakdar, and W. Hübner, *Phys. Rev. Lett.*, **100**, 116103 (2008).

⁷ M. Vogt, R. Zimmermann, and R. J. Needs, *Phys. Rev. B*, **69**, 045113 (2004).

⁸ J. Schirmer, L. S. Cederbaum, and O. Walter, *Phys. Rev. A*, **28**,

1237 (1983).

⁹ M. Altarelli, D. L. Dexter, H. M. Nussenzveig, and D. Y. Smith, *Phys. Rev. B*, **6**, 4502 (1972).

¹⁰ J. Winter, *Nucl. Phys. A*, **194**, 535 (1972).

¹¹ L. Hedin, *Phys. Rev.*, **139**, A796 (1965).

¹² G. Onida, L. Reining, and A. Rubio, *Rev. Mod. Phys.*, **74**, 601 (2002).

¹³ U. von Barth and B. Holm, *Phys. Rev. B*, **54**, 8411 (1996).

¹⁴ L. Brillouin, *J. Phys. Radium*, **32**, 373 (1932).

¹⁵ D. Pines and P. Nozières, *The Theory of Quantum Liquids* (Perseus Books Group, 1999) p. 309.

¹⁶ B. L. Altshuler, Y. Gefen, A. Kamenev, and L. S. Levitov, *Phys. Rev. Lett.*, **78**, 2803 (1997).

¹⁷ Y. Pavlyukh and J. Berakdar, *Phys. Rev. A*, **81**, 042515 (2010).